



Eurostarch

medium late, high yielding starch potato with good resistance to nematodes and diseases

Maturity	medium late	Tuber tuber shape eye depth skin flesh colour	round-oval shallow yellow, netted white
Yield	very good starch yield due high yield of tubers with high starch content		
Quality starch content	high, safe yield	Resistances Potato wart disease PCN Potato Virus Y (PVY) Potato Leafroll Virus leaf blight Blackleg Rhizoctonia common scab	D1 Ro 1, Ro 4 very high very high high high high high
Youth development	rapid, quick develop- ing canopy		
Dormancy	high – very high, very good storability	Sensitivities spraing bruising mechanical damage	low high high

Eurostarch has medium demands to soil and water supply, and is high resistant to common scab.

Eurostarch has as medium tuberisation rate and is growing medium to large sized tubers. A planting density of approx. 42,000 plants/ha is recommended.

Eurostarch should be fertilized at local standard level. Usually 160 kg/ha N (including soil supplied N_{min}) and 100 kg/ha P_2O_5 and 150 kg/ha K_2O fit to demands of Eurostarch at medium soil nutrient content classes.

Eurostarch is a high performer within medium late portfolio of starch varieties.

All particulars are based on official variety trials and/or our own experience.
However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

P. O. Box 1380 - D-21303 Lüneburg - tel.: +49 4131 748005 - fax: +49 4131 7480580
email: europlant@europlant.biz - www.europlant.biz