



**EUROPLANT**


**CAPTIVA**



## **mainly firm cooking ware potato**

- ✓ light yellow flesh colour
- ✓ medium early maturity
- ✓ smooth skin and shallow eyes
- ✓ very high marketable yield

CAPTIVA - large sized and attractive tubers

| Growing recommendations  |  |   |   |           |
|--|--|---|---|-----------|
| Site requirements  | average to better soil conditions; locations susceptible to rust spot should be avoided                                    |   |   |           |
| <div>Planting</div>  | Seed material sorting<br>35/55   | Row width   |   | Tubers/ha |
|  |  | 75 cm   | 90 cm   |           |
|  |  | 32-34 cm  | 27-28 cm  | 40,400    |
|  | Planting depth   | standard  |   |           |
|  | Seed preparation   | has a good dormancy; sprout stimulation evenly before planting (7-10 days)                    |   |           |
| Seed dressing  | dressing against Rhizoctonia solani recommended  |   |   |           |
| Herbicides   | exhibits average sensitivity to Metribuzin in the pre- and post-emergence  |   |   |           |
| Foliage  | good, average leaf health  |   |   |           |
| Harvest  | ensure gentle harvest  |   |   |           |
| Fertilization incl. return of previous crop at medium supply level                                   |  |   |   |           |
| Nitrogen (N)   | medium - compared to the local standard (2/3 before planting and 1/3 later)  |   |   |           |
| Phosphorus (P <sub>2</sub> O <sub>5</sub> )  | 100 kg/ha (water soluble phosphate at planting)  |   |   |           |
| Potassium (K <sub>2</sub> O)   | 350-270 kg/ha (to ensure the rapid storage of substances; preferably sulphate potassium, up to 80 kg/ha chloride possible) |   |   |           |
| Magnesium (MgO)  | 70 kg/ha (for yield and quality assurance)   |   |   |           |
| Variety characteristics  |  |   |   |           |
| Maturity   | medium early   | Tuber characteristics<br>Eye depth<br>Flesh colour<br>Skin colour<br>Skin<br>Number of tubers | oval<br>shallow<br>light yellow<br>yellow<br>smooth<br>low - medium |           |
| Cooking type   | B - A  |   |   |           |
| Consumption quality  | mainly firm cooking; low discolouration after cooking; good flavour  |   |   |           |
| Yield  | uniform, high tuber yield with high share of marketable yield > 50 mm  | Resistances<br><br>PCN<br><br>Potato wart disease   | Ro1 (9), Ro2/3 (9), Ro4 (9), Ro5 (9)<br>1                           |           |
| Youth development  | rapid  | Sensitivities<br><br>Common scab<br>Spraing<br>Bruising<br>Mechanical damage                  | low<br>medium<br>low<br>low - medium                                |           |
| Dormancy   | good dormancy  |   |   |           |

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (06/2021)