

medium early, mainly firm cooking ware potato

- yellow flesh colour
- roundoval tuber shape
- ✓ large grading size
- high marketable yield

COLUMBIA



COLUMBIA - large sizes and attractive tubers

Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients				
Planting Display Planting Display Planting	Seed material sorting 35/55	Row width		Tubers/ha	
		75 cm	90 cm		
		32-34 cm	27-28 cn	n 40,400	
		for large seed sizes + 2 cm			
	Planting depth	standard			
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)			
	Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	no sensitivities are known	no sensitivities are known when used properly*; seek advice			
Foliage	average leaf health; timely and regular use of fungicides against late blight is advisable				
Harvest	ensure gentle harvest				
Fertilization incl. return of	previous crop at medium supply leve	l e			
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)				
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)				
Potassium (K₂O)	350-270 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)				
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)				
Variety characteristics					
Maturity	medium early	Tuber character	istics		
Cooking type	В	Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers		roundoval shallow - medium	
Consumption quality	mainly firm cooking, low discolouration after cooking, good taste			edium - high	
Yield	large grading size with high marketable yield	Resistances PCN Potato wart		o1 (7) (a)	
Youth development	rapid	Bruising low			
Dormancy	good dormancy, good storability			ery low - low	

^{*} preliminary | (a) low sensitivities

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)