



EUROPLANT


CORDULA



early, mainly firm cooking ware potato

- ✓ yellow flesh colour
- ✓ roundoval tuber shape
- ✓ smooth skin and shallow eyes
- ✓ early shell strength

CORDULA - early culinary quality with uniform packing sizes

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting 	Seed material sorting 35/50	Row width		Tubers/ha 43,000
		75 cm	90 cm	
		30-32 cm	25-27 cm	
	Planting depth	standard		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	no sensitivities are known when used properly*			
Foliage	average leaf health; timely and regular use of fungicides against late blight is advisable			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (whole amount at planting)			
Phosphorus (P₂O₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K₂O)	350-270 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	early	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	roundoval very shallow - shallow yellow yellow smooth medium - high	
Cooking type	B			
Consumption quality	mainly firm cooking; low discolouration after cooking and good consumption quality	Resistances PCN	Ro1 (9)	
Yield	high yields with uniform sorting	Sensitivities Common scab Spraing Bruising Mechanical damage	low low - medium low medium	
Youth development	rapid			
Dormancy	good dormancy			

* preliminary

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (06/2024)