


EURODELTA - medium early starch variety with broad PCN resistance and -tolerance as well as resistance against wart disease

Growing recommendations				
Site requirements	average to better soil conditions; locations susceptible to rust spot should be avoided			
<div>Planting</div> 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	standard		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	no sensitivities are known when used properly*			
Foliage	good, average leaf health; however, regular use of fungicides is recommended			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	100 kg/ha (water soluble phosphate at planting)			
Potassium (K <sub>2</sub> O)	250–190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium early	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	roundoval medium cream yellow netted medium - high	
Starch content	very high starch performance with early starch content			
Yield	early high tuber yields			
Youth development	rapid	Resistances PCN  Nematode tolerance Potato wart disease	Ro1 (9) Pa2 (8), Pa3 (8) tolerant 1, 2 (a), 6, 18	
Dormancy	good dormancy			
		Sensitivities Common scab Spraing Bruising Mechanical damage	low high medium high	

\* preliminary | (a) low sensitivities

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)