


EUROFLORA - secures a high starch performance with high and stable starch contents; good drought tolerance together with a good leaf health add the starch segment

Growing recommendations				
Site requirements		prefers average to better soil conditions with uniform supply of water and nutrients		
<div>Planting</div> 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	1-2 cm deeper		
	Seed preparation	no special preparation necessary		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides		no sensitivities are known when used properly		
Foliage		average leaf health; timely and regular use of fungicides also against Alternaria is recommended		
Harvest		ensure gentle harvest		
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)		medium - compared to the local standard (2/3 before planting and 1/3 later)		
Phosphorus (P <sub>2</sub> O <sub>5</sub> )		100 kg/ha (water soluble phosphate at planting)		
Potassium (K <sub>2</sub> O)		250-190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)		
Magnesium (MgO)		70 kg/ha (for yield and quality assurance)		
Variety characteristics				
Maturity		medium late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Number of tubers	roundoval medium - deep light yellow yellow medium
Starch content		high and stable starch content		
Yield		high tuber yields		
Youth development		rapid	Resistances  PCN  Nematode tolerance Potato wart disease	Ro1 (9), Ro2/3 (9) Pa2 (9), Pa3 (8) tolerant 1
Dormancy		good dormancy		
			Sensitivities  Common scab Spraing Bruising Mechanical damage	very low - low medium - high low - medium low - medium

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)