

EUROLIKA - medium late starch variety with full nematode resistance

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38.500
	80 g	39 cm	33 cm	34.100
	Planting depth	standard		
	Seed preparation	germinating variety; keep cool until seed preparation		
	Seed dressing	dressing against Rhizoctonia solani is recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides also against Alternaria is recommended			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	high - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	250-190 (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin Number of tubers		round - oval medium white netted medium - high
Starch content	high starch content			
Yield	high tuber yields			
Youth development	rapid	Resistances		Ro1 (9), Ro2/3 (9), Ro5(9) Pa2 (9), Pa3 (9) 1, 2 (a), 6 (a)
Dormancy	germinating variety	PCN Potato wart disease		
		Sensitivities Common scab Spraing Bruising Mechanical damage		medium medium medium - high medium - high

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (05/2026)