

EUROMIRA - medium late starch potato with high starch yields and medium sized tubers

Growing recommendations				
Site requirements		prefers average to better soil conditions with uniform supply of water and nutrients		
Planting 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	standard		
	Seed preparation	no special preparation necessary		
	Seed dressing	dressing against Rhizoctonia solani is recommended		
Herbicides		no sensitivities are known when used properly		
Foliage		average leaf health; timely and regular use of fungicides also against Alternaria is recommended		
Harvest		ensure gentle harvest		
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)		medium - compared to the local standard (1/2 before planting and 1/2 later)		
Phosphorus (P ₂ O ₅)		100 kg/ha (water soluble phosphate at planting)		
Potassium (K ₂ O)		250-190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)		
Magnesium (MgO)		70 kg/ha (for yield and quality assurance)		
Variety characteristics				
Maturity		medium late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	round medium - deep white - cream yellow netted medium
Starch content		high starch content due to high yields		
Yield		high yields with medium sized tuber grading		
Youth development		rapid	Resistances PCN Potato wart disease	Ro1 (9), Ro2/3 (8), Ro5 (8) Pa2 (9), Pa3 (9) 1, 2 (a), 6 (a) 8/18 (a)
Dormancy		good dormancy		
			Sensitivities Common scab Spraing Bruising Mechanical damage	medium - high medium - high high medium

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (05/2026)