


EUROPURA - high starch performance with broad nematode tolerance and resistance

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	standard		
	Seed preparation	germinating variety; keep cool until seed preparation		
Seed dressing	dressing against <i>Rhizoctonia solani</i> is recommended			
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides also against <i>Alternaria</i> is recommended			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	high - compared to the local standard (1/2 before planting and 1/2 later)			
Phosphorus (P₂O₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K₂O)	250-190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	roundoval shallow - medium white - cream yellow smooth - netted medium - high	
Starch content	high starch yields due to high starch content			
Yield	medium to high tuber yields			
Youth development	rapid	Resistances PCN Nematode tolerance Potato wart disease	Ro1 (9), Ro2/3 (8), Ro5 (8) Pa2 (9), Pa3 (9) tolerant 1, 2 (a)	
Dormancy	good dormancy			Sensitivities Common scab Spraing Bruising Mechanical damage

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (09/2023)