


EUROSTARCH - medium early, robust starch variety with high tolerance against secondary growth on drought locations

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
<div>Planting</div> 	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	standard		
	Seed preparation	no special preparation necessary		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides also against Alternaria is recommended			
Harvest	standard			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	250-190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	roundoval shallow - medium white yellow netted low - medium	
Starch content	very good starch yield by high starch content			
Yield	high tuber yields			
Youth development	rapid	Resistances PCN Potato wart disease	Ro1, Ro4 1, 2 (a), 6 (a)	
Dormancy	very dormant, suitable for long-term storage	Sensitivities Common scab Spraing Bruising Mechanical damage	low low - medium medium - high medium - high	

(a) low sensitivities

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)