


EUROVIVA - late variety with a long vegetation time ensures starch performance at the highest level

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
<div>Planting</div> <div></div>	Ø g/tuber	Row width		Tubers/ha
		75 cm	90 cm	
	70 g	35 cm	29 cm	38,500
	80 g	39 cm	33 cm	34,100
	Planting depth	standard		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides also against Alternaria is recommended			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	100 kg/ha (at planting, water soluble phosphate)			
Potassium (K <sub>2</sub> O)	250-190 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	late	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers		roundoval shallow - medium light yellow yellow netted medium - high
Starch content	high starch yield with high starch contents			
Yield	high tuber yields			
Youth development	rapid	Resistances PCN  Potato wart disease		Ro1 (9) Pa2 (8), Pa 3 1, 2, 6, 18, 8 (field resistance), 38**
Dormancy	good dormancy			
		Sensitivities Common scab Spraing Bruising Mechanical damage		medium - high high high medium

\*\* tested in an EU country

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (09/2023)