

very early, mainly firm cooking ware potato

- yellow flesh colour
- round-oval tuber shape
- early high marketable yields
- ✓ variety for foil cultivation and pre-germination





FILIPA - very early, mainly firm cooking and attractive ware potato

Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting Planting	Seed material sorting 35/50	Row width		Tubers/ha
		75 cm	90 cm	
		32-34 cm	27-28 cm	40,400
		pre-germination plant 2 to 4 cm closer		
	Planting depth	standard		
	Seed preparation	germinating variety; keep cool until seed preparation		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly*			
Foliage	average leaf health; timely and regular use of fungicides against late blight is required			
Harvest	timely and gentle harvesting after achieving skin stability			
Fertilization incl. return of	previous crop at medium supply level			
Nitrogen (N)	medium - compared to the local standard (whole amount at planting)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	325-245 kg/ha (to ensure the rapid storage of substances preferably sulphate potassium)			
Magnesium (MgO)	60 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	very early	Tuber characteristics	stics	
Cooking type	В	Tuber shape Eye depth	round	
Consumption quality	good taste; stable quality	Flesh colour	yellov	V
	after cooking	Skin colour Skin	yellov	
		Number of tu		-
Yield	early high marketable yields	Resistances		
		PCN	Ro1 (Ro5 (9), Ro2/3 (8), 8)
Youth development	rapid	Sensitivities		
Dormancy	germinating variety	Common scal Spraing Bruising Mechanical da	low -	low - low medium low - low

^{*} preliminary

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (08/2023)