EUROPLANT

# GLORIETTA



## very early to early, firm cooking premium ware potato

- deep yellow flesh colour
- salad shape
- smooth skin and shallow eyes
- excellent potato flavour

# GLORIETTA

GLORIETTA - nice salad shape, smooth skin with shallow eyes and excellent flavour

| **Growing recommendations** | | | | |
|---|---|---|---|---|
| **Site requirements** | prefers average to better soil conditions with uniform supply of water and nutrients | | | |
| **Planting** | **Seed material sorting 35/55** | **Row width** | | **Tubers/ha** |
| | | **75 cm** | **90 cm** | |
| | | 28-30 cm | 23-25 cm | 46,000 |
| | **Planting depth** | plant higher tuber nest 2 cm deeper if necessary (max. 17 cm soil cover) | | |
| | **Seed preparation** | has good dormancy; sensitive to sprout breaking; a temperature shock briefly before planting (3-4 days) is beneficial | | |
| | **Seed dressing** | dressing against Rhizoctonia solani recommended | | |
| **Herbicides** | no sensitivities are known when used properly | | | |
| **Foliage** | good, average leaf health | | | |
| **Harvest** | timely and gentle harvesting after achieving skin stability | | | |
| **Fertilization** incl. return of previous crop at medium supply level | | | | |
| **Nitrogen (N)** | medium - compared to the local standard (whole amount at planting) | | | |
| **Phosphorus ($P_2O_5$)** | 100 kg/ha (water soluble phosphate at planting) | | | |
| **Potassium ($K_2O$)** | 325-245 kg/ha (to ensure the rapid storage of substances; preferably sulphate potassium, up to 80 kg/ha chloride possible) | | | |
| **Magnesium (MgO)** | 60 kg/ha (for yield and quality assurance) | | | |
| **Variety characteristics** | | | | |
| **Maturity** | very early to early | **Tuber characteristics** | | |
| **Cooking type** | A | Tuber shape<br>Eye depth<br>Flesh colour<br>Skin colour<br>Skin<br>Number of tubers | | longoval<br>very shallow - shallow<br>deep yellow<br>yellow<br>smooth<br>medium |
| **Consumption quality** | firm cooking, no discolouration after cooking; excellent potato flavour | **Resistances** | | |
| | | PCN<br>Potato wart disease | | Ro1 (9), Ro4 (9)<br>1 |
| **Yield** | medium yield with high amount of packing sizes | **Sensitivities** | | |
| **Youth development** | initially hesitant, later rapid | Common scab<br>Spraing<br>Bruising<br>Mechanical damage | | low<br>medium - high<br>low<br>low |
| **Dormancy** | good dormancy | | | |

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)

EUROPLANT Pflanzenzucht GmbH | Wulf-Werum-Str. 1 | 21337 Lüneburg GERMANY |
Tel. +49 4131 7480 05 | Fax +49 4131 7480 580 | info@europlant.biz | www.europlant.biz