EUROPLANT

# JELLY



## medium late, mainly firm cooking ware potato

- yellow flesh colour
- attractive tubers with stable shape
- very high marketable yield
- perfect adaptability to all climate conditions

# JELLY

JELLY - combines very high market yields, uniform medium to large grading size, good drought tolerance and resistance to secondary growth with good culinary quality

| **Growing recommendations** | | | | |
|---|---|---|---|---|
| **Site requirements** | no special requirements in terms of location | | | |
| **Planting** | **Seed material sorting 35/55** | **Row width** | | **Tubers/ha** |
| | | **75 cm** | **90 cm** | |
| | | 28-30 cm | 23-25 cm | 46,000 |
| | | for large tuber sizes + 4 cm | | |
| | **Planting depth** | standard | | |
| | **Seed preparation** | has a good dormancy; sprout stimulation evenly before planting (7-10 days) | | |
| | **Seed dressing** | dressing against Rhizoctonia solani recommended | | |
| **Herbicides** | exhibits average sensitivity to Metribuzin in the post-emergence | | | |
| **Foliage** | average leaf health; timely and regular use of fungicides against late blight is advisable | | | |
| **Harvest** | ensure gentle harvest | | | |
| **Fertilization** incl. return of previous crop at medium supply level | | | | |
| **Nitrogen (N)** | low - compared to the local standard (2/3 before planting and 1/3 later) | | | |
| **Phosphorus ($P_2O_5$)** | 100 kg/ha (water soluble phosphate at planting) | | | |
| **Potassium ($K_2O$)** | 350-270 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses) | | | |
| **Magnesium (MgO)** | 70 kg/ha (for yield and quality assurance) | | | |
| **Variety characteristics** | | | | |
| **Maturity** | medium late | **Tuber characteristics** | | |
| **Cooking type** | B | Tuber shape | oval | |
| | | Eye depth | shallow | |
| | | Flesh colour | yellow | |
| | | Skin colour | yellow | |
| | | Skin | smooth - netted | |
| | | Number of tubers | medium | |
| **Consumption quality** | mainly firm cooking, high consumption quality and low discolouration after cooking, suitability for processing to French fries | **Resistances** | | |
| | | PCN | Ro1 (9), Ro2/3 (9), Ro4, Ro5 | |
| | | Potato wart disease | 1, 2 (a), 6 (a) | |
| **Yield** | very high marketable yield, medium to large sized tubers, uniform grading | **Sensitivities** | | |
| | | Common scab | low | |
| | | Spraing | low | |
| | | Bruising | low | |
| **Youth development** | rapid | Mechanical damage | low - medium | |
| **Dormancy** | good dormancy | | | |

(a) low sensitivities

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (05/2026)

EUROPLANT Pflanzenzucht GmbH | Wulf-Werum-Str. 1 | 21337 Lüneburg GERMANY |
Tel. +49 4131 7480 05 | Fax +49 4131 7480 580 | info@europlant.biz | www.europlant.biz