





early, mainly firm cooking ware potato

- ✓ yellow flesh colour
- early high marketable yield
- smooth skin, shallow eyes
- early good washability

JULINKA



JULINKA - washed, brushed or fresh, always a good solution

Growing recommendat	ions			
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		28-30 cm	23-25 cm	46,000
	Planting depth	planted 2 cm deeper for high-yielding locations (max. 17 cm soil cover)		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides against late blight is required			
Harvest	ensure gentle harvest			
Fertilization incl. return of	previous crop at medium supply level			
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	350-270 kg/ha (up to 150 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	early	Tuber characteristic	cs	
Cooking type	В	Tuber shape Eye depth	oval shallo	
Consumption quality	mainly firm cooking; very stable in colour	Flesh colour	yellow	
		Skin colour Skin	yellow	
		Number of tube		
Yield	medium to high yield in a	Resistances		
	tight packing grading	PCN	Ro1 (9	9), Ro4 (9)
Youth development	initially hesitant; later rapid	Sensitivities		
Dormancy	good dormancy	Common scab Spraing	low	
		Bruising	low	
		Mechanical dam	age low - r	nedium

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)