



EUROPLANT


JULINKA



early, mainly firm cooking ware potato

- ✓ yellow flesh colour
- ✓ early high marketable yield
- ✓ smooth skin, shallow eyes
- ✓ early good washability

JULINKA - washed, brushed or fresh, always a good solution

Growing recommendations				
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting 	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		28-30 cm	23-25 cm	46,000
	Planting depth	planted 2 cm deeper for high-yielding locations (max. 1.7 cm soil cover)		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	average leaf health; timely and regular use of fungicides against late blight is required			
Harvest	ensure gentle harvest			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P₂O₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K₂O)	350-270 kg/ha (up to 150 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	early	Tuber characteristics	oval shallow yellow yellow smooth medium	
Cooking type	B			
Consumption quality	mainly firm cooking; very stable in colour			
Yield	medium to high yield in a tight packing grading	Resistances	PCN Ro1 (9), Ro4 (9)	
Youth development	initially hesitant; later rapid	Sensitivities	Common scab low Spraing low Bruising low Mechanical damage low - medium	
Dormancy	good dormancy			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)