

## early, mainly firm cooking ware potato

- ✓ yellow flesh colour
- nice, attractive tuber shape
- ✓ good culinary quality
- excellent suitability for pre-packing





## MARABEL - attractive tubers with good taste for pre-packing

Site requirements	average to better soil cond	itions; locations susceptible	to rust spot	should be avoide	
Planting	Seed material sorting	Row width		Tubers/ha	
	35/55	75 cm	90 cm		
			25-27 cm	43,000	
	Planting depth	standard			
	Seed preparation	has a good dormancy; sprout stimulation evenly befor planting (7-10 days)			
	Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	exhibits average sensitivity	ge sensitivity to Metribuzin in the post-emergence			
Foliage	good, average leaf health				
Harvest	standard				
Fertilization incl. return of	previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)				
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	100 kg/ha (water soluble phosphate at planting)				
Potassium (K <sub>2</sub> O)	350-270 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)				
Magnesium (MgO)	70 kg/ha (for yield and qua	lity assurance)			
Variety characteristics					
Maturity	early	Tuber characteristics			
Cooking type	В	Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	oval shallow yellow yellow smooth medium		
Consumption quality	mainly firm cooking; no discolouration after cooking; suitable for peeling and convenience	Resistances PCN	Ro1, Ro	o4	
Yield	products  high tuber yield with uniform large sized grading	Sensitivities  Common scab Spraing Bruising Mechanical damag	low - m	low - medium low - medium low medium	
Youth development	rapid				
Dormancy	good dormancy				

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)