

medium early, mainly firm cooking ware potato

- yellow flesh colour
- oval, attractive tuber shape
- broad potato wart resistance
- 🧹 good storability

ΟΤΟLΙΑ



OTOLIA - culinary quality with healthy foliage and extensive potato wart resistance

Growing recommendati	ons			
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		30-32 cm 25-27 cm 43,000		
		for smaller packing size -2-4 cm		
	Planting depth	standard		
	Seed preparation	no special preparation necessary		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly			
Foliage	very good leaf health (late blight), slightly senstive to early blight			
Harvest	ensure gentle harvest			
Fertilization incl. return of	previous crop at medium supply level			
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	350-270 kg/ha (up to 150 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium early	Tuber characteristics		
Cooking type	В	Tuber shape Eye depth	oval	v - medium
Consumption quality	mainly firm cooking; very low discolouration after cooking	Flesh colour Skin colour Skin Number of tubers	yellow yellow netted	
Yield	medium yields with uniform packing grading	Resistances PCN Potato wart disea), Ro2/3 (8) , 18
Youth development	initially hesitant, later rapid	Sensitivities Common scab		
Dormancy	good dormancy	Spraing Bruising Mechanical dama	low mediu low ge mediu	

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)