

very early, red skinned, mainly firm cooking ware potato

- light yellow flesh colour
- early high marketable yield
- ✓ smooth skin with shallow eyes
- ✓ very tasteful





SANIBEL - completes the very early, red skin variety segment with a very attractive, large tuber shape

Growing recommendations					
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients				
Planting	Seed material sorting 35/55	Row width			Tubers/ha
		75 cm	90 c	m	
		30-32 cm	25-27	cm	43,000
	Planting depth	standard			
	Seed preparation	germinating variety; keep cool until seed preparation			
	Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	is sensitive to Metribuzin in the pre- and post-emergence; avoid use				
Foliage	average leaf health; timely and regular use of fungicides against late blight is required				
Harvest	ensure gentle harvest				
Fertilization incl. return of previous crop at medium supply level					
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)				
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)				
Potassium (K ₂ O)	325-245 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)				
Magnesium (MgO)	60 kg/ha (for yield and quality assurance)				
Variety characteristics					
Maturity	very early	Tuber characteristics			
Cooking type	В	Tuber shape Eye depth		oval shallow	
Consumption quality	mainly firm cooking; very low discolouration after cooking	Flesh colour Skin colour Skin Number of tubers		light yello red smooth medium	ow
Yield	high; early large size tuber yield	Resistances PCN Potato wart d	isease	Ro1 (9) 1	
Youth development	rapid	Sensitivities			
Dormancy	germinating variety	Common scal Spraing Bruising Mechanical da		very low medium very low low - med	- low

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)