

medium early, red skinned, mainly firm cooking ware potato

- yellow flesh colour
- attractive tubers
- high yields
- good storability





SPINELA - red skinned, mainly firm cooking ware potato with healthy foliage and suitability for packing

a: · ·				
Site requirements	average to better soil conditions; locations susceptible to scab should be avoided			
Planting Division of the control of	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		28-30 cm	23-25 cm	46,000
		for large tuber sizes		
	Planting depth	standard		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	is sensitive to Metribuzin in the pre- and post-emergence; avoid use			
Foliage	good, average leaf health; however, regular use of fungicides is recommended			
Harvest	timely and gentle harvesting after achieving skin stability			
Fertilization incl. return of	previous crop at medium supply leve	l .		
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	350-270 kg/ha (preferably as sulphate; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics		_		
Maturity	medium early	Tuber characteris	tics	
Cooking type	В	Tuber shape Eye depth	oval	oval very shallow - shallow yellow red smooth - netted medium
Consumption quality	good culinary quality	Flesh colour Skin colour Skin Number of tubers	yellow red smoot	
Yield	medium to high yields, large sized tubers	Resistances PCN	Ro1 (9	9)
Youth development	hesitant	Sensitivities		
Dormancy	good dormancy	Common scab Spraing Bruising Mechanical da	low low - i	m - high medium m - high

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (06/2021)