

TAORMINA



medium early, mainly firm cooking export variety

- light yellow flesh skin
- very resistant against late blight
- good drought and heat tolerance
- high marketable yields





TAORMINA - high yield potential combined with high leaf health

Growing recommendat	ions			
Site requirements	prefers average to better soil conditions with uniform supply of water and nutrients			
Planting	Seed material sorting 35/50	Row width		Tubers/ha
		75 cm	90 cm	
		32-34 cm 27-28 cm 40,400		
		plant 2 to 4 cm closer for pac		
	Planting depth	standard		
	Seed preparation	has good dormancy; sensitive to sprout breaking; a temperature shock briefly before planting (3-4 days) is beneficial		
	Seed dressing	dressing against Rhizoctonia solani recommended		
Herbicides	no sensitivities are known when used properly*			
Foliage	very good leaf health (late blight)			
Harvest	standard			
Fertilization incl. return of	previous crop at medium supply level			
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	350-270 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium early	Tuber characterist	ics	
Cooking type	В	Tuber shape Eye depth	oval	oval shallow light yellow smooth - netted medium - high
Consumption quality	low discolouration after cooking	Flesh colour Skin Number of tubers	light y smoot	
Yield	high yields; large grading	Resistances		
Youth development	rapid	PCN Potato wart dis		9), Ro2/3 (8) .8
Dormancy	good dormancy	Sensitivities Common scab Spraing Bruising Mechanical dar		ow - low nedium

* preliminary

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)