



EUROPLANT


TORENIA



medium early, firm cooking ware potato

- ✓ yellow flesh colour
- ✓ very high yield
- ✓ Low Input variety
- ✓ attractive tuber shape

TORENIA - heat and drought tolerance ensure high quality and high yield

Growing recommendations				
Site requirements	average to better soil conditions; locations susceptible to rust spot should be avoided			
<div>Planting</div> <div></div>	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		30-32 cm	25-27 cm	
	Planting depth	standard		
	Seed preparation	has a good dormancy; sprout stimulation evenly before planting (7-10 days)		
Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	no sensitivities are known when used properly			
Foliage	good, average leaf health; however, regular use of fungicides is recommended			
Harvest	standard			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	low - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P ₂ O ₅)	100 kg/ha (water soluble phosphate at planting)			
Potassium (K ₂ O)	350-270 kg/ha (up to 150 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	70 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	medium early	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	longoval shallow yellow yellow netted high	
Cooking type	A - B			
Consumption quality	premium quality ware potato; low discolouration after cooking			
Yield	high yield with high market share	Resistances PCN Potato wart disease	Ro1 (9), Ro4 (9) 1	
Youth development	rapid	Sensitivities Common scab Spraing Bruising Mechanical damage	low medium - high very low - low low - medium	
Dormancy	good dormancy			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (06/2021)