



EUROPLANT


VINETA



## **early, mainly firm cooking to floury ware potato**

- ✓ light yellow flesh colour
- ✓ robust ware potato
- ✓ powerful taste
- ✓ high drought tolerance

VINETA - convinces by its robustness, early tuber development and good dryness tolerance

Growing recommendations				
Site requirements	no special requirements in terms of location			
<div>Planting</div> 	Seed material sorting 35/55	Row width		Tubers/ha
		75 cm	90 cm	
		28-30 cm	23-25 cm	46,000
	large sized lots + 2-4 cm			
	Planting depth	standard		
	Seed preparation	no special preparation necessary		
Seed dressing	dressing against Rhizoctonia solani recommended			
Herbicides	is sensitive to Metribuzin in the post-emergence; avoid use			
Foliage	good, average leaf health; however, regular use of fungicides is recommended			
Harvest	standard			
Fertilization incl. return of previous crop at medium supply level				
Nitrogen (N)	medium - compared to the local standard (2/3 before planting and 1/3 later)			
Phosphorus (P <sub>2</sub> O <sub>5</sub> )	100 kg/ha (water soluble phosphate at planting)			
Potassium (K <sub>2</sub> O)	325-245 kg/ha (up to 80 kg/ha chloride possible; on lighter soils split doses)			
Magnesium (MgO)	60 kg/ha (for yield and quality assurance)			
Variety characteristics				
Maturity	early	Tuber characteristics Tuber shape Eye depth Flesh colour Skin colour Skin Number of tubers	roundoval shallow - medium light yellow yellow netted medium - high	
Cooking type	B			
Consumption quality	mainly firm to floury cooking; strong potato flavour			
Yield	medium yields; large sized tuber grading	Resistances PCN Potato wart disease	Ro1 1	
Youth development	rapid	Sensitivities Common scab Spraing Bruising Mechanical damage	low low - medium low low - medium	
Dormancy	good dormancy			

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details. (07/2023)