


varieties, competence & service

Anuschka



very early salad potato with high marketable yield

- very attractive skin and shallow eyes
- pure yellow flesh colour
- excellent table quality
- high marketable yield

Variety characteristics			
Maturity	very early	Tuber Tuber shape Eye depth Skin Flesh colour	round-oval very shallow yellow, very smooth yellow - deep yellow
Cooking Type (EAPR)	A-B (salad type)		
Consumption quality	quality salad potato; good culinary quality; pure and stable yellow flesh colour also after boiling		
Yield	high marketable yield; medium to large sized tubers; uniform grading; suitable for early packing	Resistances PCN Potato Virus Y(PVY) Potato Leafroll Virus Leaf blight Black leg Rhizoctonia Common scab	Ro 1, Ro 4 medium - high high high medium - high medium - high
Youth development	rapid		
Dormancy	high; good storability	Sensitivities Spraing Bruising Mechanical damage	medium-low very low medium
Growing recommendations			
Site requirements	preferably should be grown on soils in good agricultural condition. Soils with high risk of spraing should be avoided. High resistance to secondary growth.		
Planting	planting density	approx. 38.000 plants/ha (seed size 35/55 mm) => check QR Code for more details	
	planting depth	normal compared with local standard	
	seed preparation	careful induction of sprouting before planting is recommended (3 - 4 days) in order to accelerate maturity, skin set and harvest ability – sensitive to broken sprouts	
	seed dressing	normal	
Foliage	routine protective control of late blight is recommended		
Herbicides	no sensitivity known		
Fertilization		further information	
Nitrogen	medium - compared with local standard		
Phosphorus	100 kg/ha (at planting, water soluble phosphate)		
Potassium	200 - 250 kg/ha		
Magnesium	60 kg/ha		