

varieties, competence & service

Constantina




medium early, good sized ware potato

- light yellow flesh colour
- smooth skin
- potato wart resistant

Constantina

medium early, good sized ware potato
with broad potato wart resistance

Variety characteristics		
Maturity	medium early	Tuber Tuber shape Eye depth Skin Flesh colour Raw discolouration
Cooking Type (EAPR)	B	
Consumption quality	light yellow flesh colour; mainly firm cooking	
Yield	medium to high yields with big sized tubers	Resistances Potato wart disease PCN Potato Leafroll Virus Leaf blight Tuber blight Rhizoctonia Common scab
Youth development	rapid	
Dormancy	high; excellent suitable for storage	Sensitivities Spraying Bruising Mechanical damage
		oval shallow yellow, slightly netted light yellow low
		NL1 (10), 2, 18 (10) Ro 1 - 3 (all 9) high high - medium high high high
		very low very low - low low - medium
Growing recommendations		
Site requirements	medium demands to the soil with a regular water supply	
Planting	planting density	approx. 43.000 plants/ha (seed size 35/55 mm) => check QR Code for more details
	planting depth	normal
	seed preparation	sprouting stimulation is generally advantageous
	seed dressing	preventive control of <i>Rhizoctonia solani</i> is generally recommended
Foliage	routine protective control of late blight is recommended	
Herbicides	no sensitivities known	
Fertilization		further information
Nitrogen	medium - compared with local standard	
Phosphorus	100 kg/ha (at planting, water soluble phosphate)	
Potassium	250 - 300 kg/ha	
Magnesium	60 kg/ha	

All particulars are based on official variety trials and/or our own experience.
However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 748005 · fax: +49 4131 7480580
mail: europlant@europlant.biz · www.europlant.biz