



Europrima

early crisp variety

| | | | |
|--|---|---|---|
| Maturity | early | Tuber Tuber shape Eye depth Flesh colour Skin | round - short-oval shallow - medium light yellow yellow, slightly netted |
| Yield Marketable yield | high high - very high, uniform large sized grading | | |
| Quality Dry matter Reducing sugars Crisp colour Raw discolouration | medium very low very good very low - low | Resistances PCN Potato wart disease Virus diseases Leaf blight Blackleg Common scab | Ro 1, Ro 4 D1 very high high high medium - high |
| | Youth development | | |
| Dormancy | high, good storability with very low sugar content | Sensitivities Spraying Bruising Mechanical damage | low medium medium |

Europrima has an early and high yield of uniform large sized tubers. The tuberization rate is medium to low, a plant density of approx. 42,000 plants/ha is recommended for early harvesting. For late scheduled harvesting date approx. 46,000 plants/ha are recommended. Careful inducing of sprouting before planting is beneficial for quick emergence and rapid development.

Europrima should be harvested at progressed maturity in order to ensure excellent processing quality.

Europrima has very good storability. The content of reducing sugars is very low and the processing quality is excellent, even after storage.

All particulars are based on official variety trials and/or our own experience.
However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 7480-05 · fax: +49 4131 7480-580
mail: europlant@europlant.biz · www.europlant.biz