



# Eurostarch

medium late, high yielding starch potato with good resistance to nematodes and diseases

Maturity	medium late	Tuber Tuber shape Eye depth Skin Flesh colour	round-oval shallow yellow, netted white
Yield	very good starch yield due high yield of tubers with high starch content		
Quality Starch content	very high, safe yield	Resistances Potato wart disease PCN Potato Virus Y (PVY) Potato Leafroll Virus Leaf blight Blackleg Rhizoctonia Common scab	D 1 Ro 1, Ro 4 very high very high high high high high
Youth development	rapid, quick developing canopy		
Dormancy	high - very high, very good storability		
		Sensitivities Spraing Bruising Mechanical damage	low high high

**Eurostarch** has medium demands to soil and water supply, and is high resistant to common scab.

**Eurostarch** has a medium tuberisation rate and is growing medium to large sized tubers. A planting density of approx. 42,000 plants/ha is recommended.

**Eurostarch** should be fertilized at local standard level. Usually 160 kg/ha N (including soil supplied  $N_{min}$ ) and 100 kg/ha  $P_2O_5$  and 150 kg/ha  $K_2O$  fit to demands of **Eurostarch** at medium soil nutrient content classes.

**Eurostarch** is a high performer within medium late portfolio of starch varieties.

All particulars are based on official variety trials and/or our own experience.  
However, as potatoes are natural products we cannot undertake any liability for these details.

## EUROPLANT Pflanzenzucht GmbH

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 7480-05 · fax: +49 4131 7480-580  
mail: [europlant@europlant.biz](mailto:europlant@europlant.biz) · [www.europlant.biz](http://www.europlant.biz)