


varieties, competence & service

# Gunda



early, floury cooking quality ware potato

- nice tuber shape
- excellent table quality
- suitable for washing and pre-packing

Variety characteristics		
<b>Maturity</b>	early	<b>Tuber</b>  Tuber shape Eye depth Skin Flesh colour Raw discolouration
<b>Cooking Type (EAPR)</b>	B-C	
<b>Consumption quality</b>	mealy cooking quality ware potato, very good taste, clear yellow flesh colour, very low discolouration after cooking	
<b>Yield</b>	high marketable yield, medium sized tubers, very uniform grading, very good suitability for packing	<b>Resistances</b>  Potato wart disease PCN Potato Leafroll Virus Leaf blight Tuber blight Rhizoctonia Common scab
<b>Youth development</b>	initially hesitant, later rapid	
<b>Dormancy</b>	high, excellent suitable for storage	<b>Sensitivities</b>  Spraing Bruising Mechanical damage
<b>Growing recommendations</b>		
<b>Site requirements</b>	medium demands to the soil quality and water supply	
<b>Planting</b>	planting density	approx. 39.000 plants/ha (seed size 35/55 mm) => check QR Code for more details
	planting depth	normal compared with local standard
	seed preparation	sprouting stimulation 3 days before planting
	seed treatment	preventive control of <i>Rhizoctonia solani</i> is generally recommended
<b>Foliage</b>	routine protective control of early blight (Alternaria) is recommended	
<b>Herbicides</b>	no sensitivities known	
<b>Fertilization</b>		<b>further information</b>
<b>Nitrogen</b>	medium - compared with local standard	
<b>Phosphorus</b>	100 kg/ha (at planting, water soluble phosphate)	
<b>Potassium</b>	200 - 250 kg/ha	
<b>Magnesium</b>	60 kg/ha	

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.