


varieties, competence & service

Theresa



medium early quality ware potato

- light yellow flesh colour
- cooking type C
- nice shape and uniform tubers

Variety characteristics			
Maturity	medium early	Tuber Tuber shape Eye depth Skin Flesh colour	roundoval shallow - medium lightly netted light yellow
Cooking Type (EAPR)	C		
Consumption quality	light yellow flesh colour, very low discolouration after cooking, good taste		
Yield	medium	Resistances PCN Potato Virus Y (PVY) Leaf blight Tuber blight Black leg Rhizoctonia Common scab	Ro 1(9), 4(9) medium - high medium high - high high high high - very high
Youth development	rapid		
Dormancy	high, good storability	Sensitivities Spraing Bruising Mechanical damage	very low low medium
Growing recommendations			
Site requirements	preferably grown on soils with uniform water supply		
Planting	planting density	approx. 46,000 plants/ha (seed size 35/55 mm) => check QR Code for more details	
	planting depth	normal	
	seed preparation	sprouting stimulation before planting	
	seed dressing	preventive control of <i>Rhizoctonia</i> is generally recommended	
Foliage	Theresa has a good foliage, routine protective control of late blight is recommended		
Herbicides	no sensitivities known		
Fertilization			further information
Nitrogen	medium - compared to local standard		
Phosphorus	100 kg/ha (at planting, water soluble phosphate)		
Potassium	200 - 250 kg/ha		
Magnesium	60 kg/ha		

All particulars are based on official variety trials and/or our own experience.
However, as potatoes are natural products we cannot undertake any liability for these details.