


Mainly firm cooking variety with broad PCN-resistance and tolerance

Variety characteristics			
Maturity	medium early	Tuber Tuber shape Eye depth Skin Flesh colour	oval shallow yellow, smooth yellow - deep yellow
Cooking Type (EAPR)	B		
Consumption quality	yellow flesh colour; cooking type B; very low discoloration after cooking		
Yield	high yield with nice uniform packing grading	Resistances PCN PCN-tolerance Potato wart disease	Ro1, 3, 5 (9), Pa2 (7), Pa3 (6)* High 1
Youth development	initially hesitant, later rapid		
Dormancy	high, good storability	Sensitivities Spraing Bruising Mechanical damage Scab	medium low medium medium - low
Growing recommendations			
Site requirements	preferably grown on medium to good soils with uniform water and nutrient supply; care should be taken on good agricultural condition of soil. Soils with high risk of spraing should be avoided		
Planting	planting density	approx. 38,000 plants / ha (seed size 35/55 mm) for 40/60 packing size 2 cm deeper than usual	
	planting depth	normal	
	seed preparation	strong dormancy; application of temperature shock 3 - 4 days before planting necessary	
	seed dressing	preventive control of Rhizoctonia solani is generally recommended	
Foliage	good, average Foliage		
Herbicides	slightly sensitive to Metribuzin, dose rates post emergence application should be avoided		
Fertilization		further information	
Nitrogen	normal local standard level (approx. 160 kg/ha, incl. soil supplied Nmin)		
Phosphorus	100 kg/ha (at planting, water soluble phosphate)		
Potassium	200 - 250 kg/ha		
Magnesium	60 kg/ha		

*) preliminary result resistance expected

All particulars are based on official variety trials and/or our own experience. However, as potatoes are natural products we cannot undertake any liability for these details.

EUROPLANT Pflanzenzucht GmbH

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 748005 · fax: +49 4131 7480580
mail: europlant@europlant.biz · www.europlant.biz