


## Nevada – for starting the French Fries season

Variety characteristics			
<b>Maturity</b>	early	<b>Tuber</b>  Tuber shape Eye depth Skin Flesh colour	roundoval shallow rough yellow - deep yellow
<b>Cooking Type (EAPR)</b>	BC		
<b>Processing quality</b>	good frying colour from the beginning		
<b>Yield</b>	medium to high yield	<b>Resistances</b>  PCN	Ro 1 (9)
<b>Youth development</b>	rapid		
<b>Dormancy</b>	high, processing suitability French Fries >8°C	<b>Sensitivities</b>  Leaf Scab Spraing Bruising Mechanical damage	medium - high low medium low medium
<b>Growing recommendations</b>			
<b>Site requirements</b>	preferably grown on medium soils with uniform water and nutrient supply		
<b>Planting</b>	planting density	approx. 38,000 plants/ha (seed size 35/55 mm)	
	planting depth	normal	
	seed preparation	temperature shock 3 days before planting	
	seed dressing	preventive control of Rhizoctonia solani is generally recommended	
<b>Foliage</b>	protective control of late blight within the usual spraying program is recommended		
<b>Herbicides</b>	under normal application programs, no sensitivities known		
Fertilization		further information	
<b>Nitrogen</b>	normal local standard level		
<b>Phosphorus</b>	100 kg/ha (at planting, water soluble phosphate)		
<b>Potassium</b>	200 - 250 kg/ha (medium soil supply class, preferably potassium sulphate)		
<b>Magnesium</b>	60 kg/ha		

All particulars are based on official variety trials and/or our own experience.  
However, as potatoes are natural products we cannot undertake any liability for these details.

**EUROPLANT Pflanzenzucht GmbH**

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 748005 · fax: +49 4131 7480580  
mail: europlant@europlant.biz · www.europlant.biz