


Variety characteristics			
<b>Maturity</b>	early to medium early	<b>Tuber</b>	
<b>Cooking Type (EAPR)</b>		Tuber shape Eye depth Skin Flesh colour	round oval shallow-medium rough light yellow
<b>Processing quality</b>	stable colour with good results, light yellow flesh colour		
<b>Yield</b>	high yield with marketable results	<b>Resistances</b>	
<b>Youth development</b>	rapid	PCN Common Scab	Ro 1, (9), 4 * medium
<b>Dormancy</b>	good storability, processing quality crisps +8°C *	<b>Sensitivities</b>	
		Spraing Bruising Mechanical damage	low low - medium low - medium
Growing recommendations			
<b>Site requirements</b>	preferably grown on medium soils with uniform water and nutrient supply. Scab endangered locations should be avoided.		
<b>Planting</b>	planting density	approx. 43.000 plants/ha (seed size 35/55 mm); => check QR Code for more details	
	planting depth	normal	
	seed preparation	temperature shock 3 days before planting	
	seed dressing	preventive control of Rhizoctonia solani is generally recommended	
<b>Foliage</b>	Protective control of late blight within the usual spraying programm is recommended		
<b>Herbicides</b>	no sensitivities known		
Fertilization		further information	
<b>Nitrogen</b>	low-medium - compared to local standard		
<b>Phosphorus</b>	100 kg/ha (at planting, water soluble phosphate)		
<b>Potassium</b>	200 - 250 kg/ha		
<b>Magnesium</b>	60 kg/ha		

\*) preliminary

All particulars are based on official variety trials and/or our own experience.  
However, as potatoes are natural products we cannot undertake any liability for these details.

**EUROPLANT Pflanzenzucht GmbH**

P. O. Box 1380 · D-21303 Lüneburg · tel.: +49 4131 748005 · fax: +49 4131 7480580  
mail: europlant@europlant.biz · www.europlant.biz